

# Strengthening respiratory protection from SARS-CoV-2 in certain general practice contexts

**Anthony D LaMontagne**, Rhea Martin, Ruby Biezen, Barbara Hunter, Mukesh Haikerwal, Karyn Alexander, Lena Sancic, Jo-Anne Manski-Nankervis

**THIS BRIEF CONTRIBUTION** addresses one element of the comprehensive infection prevention and control (IPC) programs needed to prevent work-related SARS-CoV-2 infection in healthcare workers: respiratory protection. Further, this contribution applies specifically to the general practice context.

Most IPC guidance to date has been based on assumptions that transmission of SARS-CoV-2 occurs via large droplets that travel up to 1.5–2.0 metres and by contact with contaminated surfaces, and that aerosols only play a part in transmission of the virus during aerosol-generating procedures (AGPs). While most current evidence supports these as the predominant routes of transmission,<sup>1</sup> evidence that airborne transmission of SARS-CoV-2 might occur outside of AGPs is mounting.<sup>2,3</sup> Although evidence of airborne transmission is neither conclusive nor suspected to be a major route of transmission, we recommend a precautionary approach to respiratory protection for potentially exposed healthcare workers until the efficacy of surgical masks can be proven. In particular, we propose:

*Where there is community transmission, or other reasons to expect a high probability of patients who are SARS-CoV-2 positive, particulate filter respirators such as N95/P2 should be used for throat and deep*

*nasal swab collection (SARS-CoV-2 polymerase chain reaction testing) and other close contact with potentially infected patients in general practice settings including respiratory clinics and drive-through testing facilities.*

An N95/P2 respirator is specifically designed to fit around the face for respiratory protection and to prevent the inhalation of  $\geq 95\%$  of airborne particles. A recent systematic review and meta-analysis found that N95 respirators provided significantly stronger protection from severe acute respiratory syndrome (SARS), Middle East respiratory syndrome (MERS) and SARS-CoV-2 viral transmission than surgical masks.<sup>4</sup>

Recent guidance from The Royal Australian College of General Practitioners (RACGP) recommends surgical masks/droplet precautions for throat and deep nasal swab collection.<sup>5</sup> For GP-led COVID-19 respiratory clinics,<sup>6</sup> the RACGP also recommends droplet precautions for close patient contact, and N95/P2 respirators only for contact with patients exhibiting severe pneumonia-like symptoms.

Commonwealth guidelines accessed 16 August 2020 recommend the use of an N95/P2 respirator only when performing AGP on patients with confirmed or suspected COVID-19.<sup>7</sup>

Victorian Department of Health and Human Services guidance (issued 10 September 2020) is more open to interpretation with respect to respiratory protection, recommending the use of N95/P2 respirators where patients with suspected or confirmed COVID-19 are

being ‘cohorted’ (presumably meaning co-located, as in a COVID-19 ward), where AGPs are being performed or where an ‘unplanned AGP’ might occur (eg due to ‘screaming, shouting, crying out, and vomiting’).<sup>8</sup> Such patient behaviour may be more common in institutional settings, but it could also occur in general practice contexts. In summary, our recommendation could be read as consistent with Victorian guidance, but would exceed the current RACGP and Commonwealth recommendations.

Occupational health and safety (OHS) law is also relevant. While legislation varies by state or territory, the general requirement is that employers must protect employees from recognised risks to the extent feasible. OHS law is subject to interpretation, but it is clear that there is a large and concerning number of occupational infections overall in Victoria, and this includes 36 infections (at 23 August 2020) related to general practice clinics.<sup>9</sup> It is not currently known whether or how many occupational infections – in any context – occurred via aerosol transmission. Carefully designed studies and time will be required to answer this question. In the meantime, the use of particle-filtering respirators is a clearly feasible risk-mitigation strategy that we should implement now alongside other measures.

It is crucial to emphasise that respiratory protection and appropriate fit testing is only one element in a comprehensive IPC program.<sup>10,11</sup> Other program elements include, for example, thorough IPC training for all staff, eye

protection, dedicated donning and doffing areas, physical distancing, frequent and rigorous hand sanitation, meticulous mechanical cleaning of patient care areas, and modification of staffing and/or workloads to accommodate increased demands posed by attention to IPC.

We suggest that the RACGP and Commonwealth and Victorian authorities review their guidance to consider the use of particle-filtering respirators in the specified general practice context as a precautionary approach. This would also serve to harmonise recommended best practice in this regard across these three important authorities for the general practice community.

The protection of the healthcare workforce from occupational infection is an urgent priority, and we argue that our recommendation constitutes a relatively modest, feasible and precautionary refinement of recommended best practice.

*First published online 30 September 2020.*

#### Authors

Anthony D LaMontagne ScD, MA, MEd, Professor of Work, Health & Wellbeing, Institute for Health Transformation, Faculty of Health, Deakin University, Vic. [tony.lamontagne@deakin.edu.au](mailto:tony.lamontagne@deakin.edu.au)

Rhea Martin MPH, Infection Control Consultant, Department of General Practice, Melbourne Medical School, University of Melbourne, Vic

Ruby Biezen MAppSc, PhD, Research Fellow, Department of General Practice, Melbourne Medical School, University of Melbourne, Vic

Barbara Hunter PhD, Qualitative Research Fellow, Department of General Practice, Melbourne Medical School, University of Melbourne, Vic

Mukesh Haikerwal MBChB, DipIMC RCS (Ed), DRCOG, FRACGP (Life), FAMA (Gold Medal), GP Director, Cirqit Health, Vic; Honorary Enterprise Professor, University of Melbourne, Vic

Karyn Alexander MBChB, PhD, MPH, DCH, FRACGP, Research Fellow, Department of General Practice, Melbourne Medical School, University of Melbourne, Vic

Lena Sanci MBBS, PhD, FRACGP, Professor & Department Head, Department of General Practice, Melbourne Medical School, University of Melbourne, Vic

Jo-Anne Manski-Nankervis BSc (Hons), MBBS (Hons), CHIA, PhD, FRACGP, Associate Professor – General Practice, Melbourne Medical School, University of Melbourne, Vic

Competing interests: ADL reports personal fees as a sub-contracted consultant from The University of Melbourne. RM reports funding for safety evaluation of drive through COVID-19 clinic from the Australian Government Department of Health. RB reports grants from the Australian Government Department of Health. MH reports he is a general medical practitioner and the Director of Cirqit Health, which has delivered a GP Respiratory Disease Clinic. He is also currently the Chair of Council of the Australian

Medical Association (AMA) Victoria. MH was the former Chair of the Council of the World Medical Association, and the former President of the AMA at both State and Federal levels. KA reports that she is a general medical practitioner and the Director of Cirqit Health which has delivered a GP Respiratory Disease Clinic. LS reports grants from Federal Government. JM reports funding for safety evaluation of drive through COVID-19 clinic from the Australian Government Department of Health.

Provenance and peer review: Not commissioned, peer reviewed.

#### Acknowledgements

Evaluation of the Altona North Health Hub Drive Through Testing Clinic was funded by the Australian Government Department of Health. The views and opinions expressed in this commentary are those of the authors and do not necessarily reflect those of the Department of Health.

Citation: LaMontagne AD, Martin R, Biezen R, et al. Strengthening respiratory protection from SARS-CoV-2 in certain general practice contexts. *Aust J Gen Pract* 2020;49 Suppl 38. doi: 10.31128/AJGP-COVID-38.

#### References

- Conly J, Seto WH, Pittet D, Holmes A, Chu M, Hunter PR. Use of medical face masks versus particulate respirators as a component of personal protective equipment for health care workers in the context of the COVID-19 pandemic. *Antimicrob Resist Infect Control* 2020;9:126. doi: 10.1186/s13756-020-00779-6.
- MacIntyre CR, Ananda-Rajah M, Nicholls M, Quigley AL. Current guidelines for respiratory protection of Australian health care workers against COVID-19 are not adequate and national reporting of health worker infections is required. *Med J Aust* 2020.
- Gehanno JF, Bonnetterre V, Andujar P, et al. How should data on airborne transmission of SARS-CoV-2 change occupational health guidelines? *Occup Environ Med* 2020;77:736. doi: 10.1136/oemed-2020-106707.
- Chu DK, Akl EA, Duda S, Solo K, Yaacoub S, Schünemann HJ. Physical distancing, face masks, and eye protection to prevent person-to-person transmission of SARS-CoV-2 and COVID-19: A systematic review and meta-analysis. *Lancet* 2020;395(10242):i973-87. doi: 10.1016/S0140-6736(20)31142-9.
- The Royal Australian College of General Practitioners. COVID-19 infection-control principles. East Melbourne, Vic: RACGP, 2020.
- The Royal Australian College of General Practitioners. Guiding principles for the establishment and ongoing management of GP-led COVID-19 respiratory clinics. East Melbourne, Vic: RACGP, 2020.
- Department of Health. Personal protective equipment (PPE) for the health workforce during COVID-19. Canberra, ACT: DoH, 2020.
- Department of Health and Human Services. Coronavirus disease 2019 (COVID-19): Infection prevention and control guideline. Melbourne, Vic: DHHS, 2020.
- Department of Health and Human Services. Protecting our healthcare workers. Melbourne, Vic: DHHS, 2020.
- National Health and Medical Research Council. Australian guidelines for the prevention and control of infection in healthcare. Canberra, ACT: NHMRC, 2019.
- Australasian Faculty of Occupation and Environmental Medicine. COVID-19 Workplace Risk Management. Sydney, NSW: RACP, 2020. Available at [www.racp.edu.au/docs/default-source/news-and-events/covid-19/afoem-covid-19-workplace-risk-management.pdf?sfvrsn=d035f1a\\_8](http://www.racp.edu.au/docs/default-source/news-and-events/covid-19/afoem-covid-19-workplace-risk-management.pdf?sfvrsn=d035f1a_8) [Accessed 29 September 2020].